arXiv:1503.00498v6 [math.AT] 3 Feb 2016

Occupants in manifolds

Steffen Tillmann and Michael S. Weiss

Abstract. Let K be a subset of a smooth manifold M. In some cases, functor calculus methods lead to a homotopical formula for $M \setminus K$ in terms of the spaces $M \setminus S$, where S runs through the finite subsets of K.

1. Occupants in a submanifold

1.1. Formulation of the problem. Imagine a smooth manifold M and a compact smooth submanifold L, both with empty boundary, of dimensions m and ℓ respectively. We look for a homotopical description of $M \setminus L$ in terms of the spaces $M \setminus S$, where S runs through the finite subsets of L. The finite subsets S of L could be regarded as finite sets of occupants.

For one of the more geometric formulations of the problem, choose a Riemannian metric on L. Instead of working with finite subsets S of L, we work with thickenings of finite subsets of L and we pay attention to inclusions of one such thickening in another. More precisely we work with pairs (S, ρ) where S is a finite subset of L and ρ is a function from S to the positive real numbers subject to two conditions.

- For each $s \in S$, the exponential map $\exp_s$ at s is defined and regular on the (compact) disk of radius $\rho(s)$ about the origin in $T_s L$.
- The images in L of these disks under the exponential maps $\exp_s$ are pairwise disjoint.

For a pair (S, ρ) satisfying the two conditions, let $V_L(S, \rho) \subset L$ be the union of the open balls of radius $\rho(s)$ about points $s \in S$. Then $V_L(S, \rho)$ is an open subset of L and is diffeomorphic to $\mathbb{R}^\ell \times S$. The inclusion of $M \setminus V_L(S, \rho)$ in $M \setminus S$ is a homotopy equivalence.

Let $C_k(L)$ be the space of unordered configurations of k points in L. For fixed $k \geq 0$, the pairs (S, ρ) that satisfy the two conditions and the additional condition $|S| = k$ form a space $C_k^{\mathrm{fat}}(L)$, an open subspace of the total space of some k-dimensional vector bundle on $C_k(L)$. The forgetful projection $C_k^{\mathrm{fat}}(L) \to C_k(L)$ is a fiber bundle projection with contractible fibers, hence a homotopy equivalence.

2010 *Mathematics Subject Classification.* Primary 57R19, 55P65.

This project was supported by the Humboldt foundation through a Humboldt professorship for Michael Weiss, 2012-2017.

Form the topological disjoint union of the spaces $C_k^{\mathrm{fat}}(L)$ for all $k \geq 0$ and view that as a (topological) poset $\mathcal{P}(L)$ where

$$(S, \rho) \leq (T, \sigma)$$

means simply that $V_L(S, \rho) \subset V_L(T, \sigma)$. The poset $\mathcal{P}(L)$ can also be viewed as a category. A contravariant functor Φ from $\mathcal{P}(L)$ to spaces is defined by

$$\Phi(S, \rho) = M \smallsetminus V_L(S, \rho). \tag{1.1.1}$$

There is a map

$$M \smallsetminus L \longrightarrow \operatorname{holim} \Phi\ , \tag{1.1.2}$$

determined by the inclusions $M \smallsetminus L \to \Phi(S, \rho)$ for $(S, \rho) \in \mathcal{P}(L)$; the precise meaning of holim Φ will be clarified in a moment. With a view to occupants and the problem of finding unoccupied places, we ask:

- is the map (1.1.2) a weak homotopy equivalence ?

The question has a more numerical variant. For $j \geq 0$ let $\mathcal{P}j(L)$ be the subspace and full topological sub-poset of $\mathcal{P}(L)$ consisting of all (S, ρ) in $\mathcal{P}(L)$ that satisfy $|S| \leq j$. There is a map

$$M \smallsetminus L \longrightarrow \operatorname{holim} \Phi|_{\mathcal{P}j(L)} \tag{1.1.3}$$

determined by the inclusions $M \smallsetminus L \to \Phi(S, \rho)$. We ask:

- is the map (1.1.3) highly connected ? Is there a lower bound for the connectivity, expressed in terms of j , which tends to infinity with j ?

In reading these questions, keep in mind that Φ has some continuity properties. This affects the meaning or definition of the homotopy inverse limits. Here is a quick definition using the fact that Φ is a subfunctor of a constant functor. More details can be found in definition 1.1.2 below. Let $N\mathcal{P}(L)$ be the nerve of $\mathcal{P}(L)$, a simplicial space. So $N_r\mathcal{P}(L)$ is the *space* of order-reversing maps u from $[r] = \{0, 1, \ldots, r\}$ to $\mathcal{P}(L)$. *Order-reversing* means that $u(0) \geq u(1) \geq \cdots \geq u(r)$ in $\mathcal{P}(L)$. Now holim Φ in (1.1.2) can be described as a subspace of the space of all maps from the geometric realization $|N\mathcal{P}(L)|$ to M, with the compact-open topology. A map $f\colon |N\mathcal{P}(L)| \to M$ belongs to that subspace if and only if for every r and $u \in N_r\mathcal{P}(L)$ with characteristic map $c_u\colon \Delta^r \to |N\mathcal{P}(L)|$, the composition fc_u lands in $\Phi(u(r)) \subset M$. This description gives a rather good idea what holim Φ is: the space of all homotopy coherent ways to choose a place in M when some places in L are occupied. — The homotopy limit in (1.1.3) can be defined analogously.

THEOREM 1.1.1. *If $m - \ell \geq 3$, then the map in* (1.1.2) *is a weak homotopy equivalence and the map in* (1.1.3) *is* $(1 + (j + 1)(m - \ell - 2))$*-connected.*

The proof of theorem 1.1.1 is given at the end of this section. It is based on a reduction to standard theorems in manifold calculus as found in [**11**] and [**8**]. (See also remark 1.3.5 below.) The main idea is this: apply manifold calculus to the contravariant functor F defined by $F(V) = M \smallsetminus V$ for open subsets V of L. Then the left-hand side of (1.1.2) is $F(L)$ and the right-hand side is very reminiscent of $(T_\infty F)(L)$, where $T_\infty F$ is the "Taylor series" of F in the sense of manifold calculus. Therefore the work consists mainly in showing that F is analytic. (But the definition $F(V) := M \smallsetminus V$ should be regarded as provisional.)

We begin with a slightly more systematic description or definition of the homotopy inverse limits in (1.1.2) and (1.1.3). The idea is to use the trusted formula

of Bousfield-Kan [**3**] while paying attention to topologies where appropriate. Let $\Gamma_r(\Phi)$ be the space of (continuous) sections, with the compact-open topology, of the fiber bundle $E^!_r(\Phi) \to N_r\mathcal{P}(L)$ such that the fiber over a point $u \in N_r\mathcal{P}(L)$ is $\Phi(u(r))$. So if u is given by a string $(S_0, \rho_0) \geq \cdots \geq (S_r, \rho_r)$ in $\mathcal{P}(L)$, then the fiber over u is $\Phi(S_r, \rho_r) = M \smallsetminus V_L(S_r, \rho_r)$.

Definition 1.1.2. The homotopy limit in (1.1.2) is $\mathrm{Tot}\,([r] \mapsto \Gamma_r(\Phi))$.

Remark 1.1.3. The cosimplicial space $[r] \mapsto \Gamma_r(\Phi)$ is Reedy fibrant. This means that for each $r \geq 0$ the *matching map*

$$\Gamma_r(\Phi) \longrightarrow \lim_{[r]\to[q]} \Gamma_q(\Phi)$$

determined by the non-identity (co)degeneracy operators is a fibration. (Here $q < r$ and $[r] \to [q]$ stands for a monotone surjection; see e.g. [**5**] for more details.) It is a desirable property to have because a map $X \to Y$ between Reedy fibrant cosimplicial spaces which is a degreewise weak equivalence induces a weak equivalence $\mathrm{Tot}(X) \to \mathrm{Tot}(Y)$. Sketch of a proof showing that $[r] \mapsto \Gamma_r(\Phi)$ is Reedy fibrant: since $E^!_r(\Phi) \to N_r\mathcal{P}(L)$ is a fiber bundle it is enough to note that the simplicial space $[r] \mapsto N_r\mathcal{P}(L)$ is Reedy cofibrant. (We use HELP, the homotopy extension lifting property.) It is Reedy cofibrant because, for every $r \geq 0$, the latching map

$$\operatorname*{colim}_{[r]\to[q]} N_q\mathcal{P}(L) \longrightarrow N_r\mathcal{P}(L)$$

(where $q < r$ etc.) is the inclusion of one ENR, Euclidean neighborhood retract, in another ENR as a closed subspace. Such an inclusion is a cofibration [**9**, ch.III,Thm.3.2].

As a corollary of this observation, the homotopy type of holim Φ is independent of the choice of Riemannian metric on L. If we choose two distinct Riemannian metrics μ and μ' on L and call the corresponding functors Φ and Φ' , then there is a smooth path of Riemannian metrics from μ to μ'. Using that it is easy to produce a zigzag of degreewise weak equivalences between Reedy fibrant cosimplicial spaces starting with the cosimplicial space $[r] \mapsto \Gamma_r(\Phi)$ and ending with $[r] \mapsto \Gamma_r(\Phi')$. This leads to a zigzag of weak equivalences relating holim Φ to holim Φ'. For descriptions of (something weakly equivalent to) holim Φ which do not rely on any choice of Riemannian metric on L, see [**13**] which is a sequel to this paper.

1.2. Discrete variants. Write $\delta\mathcal{P}(L)$ for the discrete variant of $\mathcal{P}(L)$. So $\delta\mathcal{P}(L)$ is a discrete poset and there is a map of posets $\delta\mathcal{P}(L) \to \mathcal{P}(L)$ which is bijective and *full*. That is, $(S, \rho) \leq (T, \sigma)$ has the same meaning in $\delta\mathcal{P}(L)$ and in $\mathcal{P}(L)$. That map from $\delta\mathcal{P}(L)$ to $\mathcal{P}(L)$ induces a map of homotopy inverse limits:

$$\operatorname*{holim}_{(S,\rho)\text{ in }\mathcal{P}(L)} \Phi(S,\rho) \longrightarrow \operatorname*{holim}_{(S,\rho)\text{ in }\delta\mathcal{P}(L)} \Phi(S,\rho) \tag{1.2.1}$$

Similarly, restricting cardinalities of configurations we have a comparison map

$$\operatorname*{holim}_{(S,\rho)\text{ in }\mathcal{P}j(L)} \Phi(S,\rho) \longrightarrow \operatorname*{holim}_{(S,\rho)\text{ in }\delta\mathcal{P}j(L)} \Phi(S,\rho) \tag{1.2.2}$$

LEMMA 1.2.1. *The maps* (1.2.1) *and* (1.2.2) *are weak equivalences.*

PROOF. We prove that the map (1.2.1) is a weak equivalence; the other statement has a similar proof. — The first idea is to replace the topological poset $\mathcal{P}(L)$ by a simplicial poset $[t] \mapsto P_t$. Therefore let P_t be the set (alias discrete space) of continuous maps from Δ^t to the underlying space of $\mathcal{P}(L)$. For $\sigma, \tau \in P_t$ we write $\sigma \le \tau$ to mean that $\sigma(x) \le \tau(x)$ for all $x \in \Delta^t$. In particular the poset P_0 is identified with $\delta\mathcal{P}(L)$.

For a morphism in Δ alias monotone map $\alpha\colon [t] \to [u]$, let Φ_α be the contravariant functor from $P_u \times \Delta^t$ to spaces obtained by composing

$$P_u \times \Delta^t \xrightarrow{\mathrm{id}\times\alpha_*} P_u \times \Delta^u \xrightarrow{\text{evaluation}} \mathcal{P}(L)$$

with Φ. Here Δ^t and Δ^u are regarded as topological posets with a trivial ordering but with a nontrivial (standard) topology. In particular, Φ_α is $\Phi|_{\delta\mathcal{P}(L)}$ when α is the identity map of the object $[0]$ in Δ. Since Φ_α is a continuous functor, we need to be explicit about holim Φ_α if we want to use it. It is defined as

$$\mathrm{Tot}\Big([r] \mapsto \prod_{\sigma_0 \ge \cdots \ge \sigma_r \ \in P_u} \lim(\Phi \circ \sigma_r\alpha_*)\Big)$$

where $\alpha_*\colon \Delta^t \to \Delta^u$ is the map covariantly induced by α and $\lim(\Phi \circ \sigma_r\alpha_*)$ is short for the space of maps $\Delta^t \to M$ which take $x \in \Delta^t$ to an element of

$$\Phi(\sigma_r(\alpha_*(x))) = M \smallsetminus V_L(\sigma_r(\alpha_*(x))) \subset M\,.$$

Now $\alpha \mapsto$ holim Φ_α is a contravariant functor from the twisted arrow category $\mathrm{tw}(\Delta)$ of Δ to the category of spaces. (The objects of the twisted arrow category $\mathrm{tw}(\mathcal{D})$ of a small category $\mathcal{D}$ are morphisms $f\colon c \to d$ in $\mathcal{D}$, and a morphism from $f\colon c_0 \to d_0$ to $g\colon c_1 \to d_1$ is a pair of morphisms $h\colon c_0 \to c_1$ and $k\colon d_1 \to d_0$ such that $f = kgh$.) The map (1.2.1) can be obtained by composing the arrows in the lower row of the commutative diagram

(1.2.3)
$$\begin{array}{ccccc} & & \lim\limits_{\alpha \text{ in } \mathrm{tw}(\Delta)} \text{holim } \Phi_\alpha & \xrightarrow{\text{incl.}} & \prod_\alpha \text{holim } \Phi_\alpha \\ & \nearrow & \big\downarrow \text{incl.} & & \big\downarrow \text{proj.} \\ \text{holim } \Phi & \longrightarrow & \lim\limits_{\alpha \text{ in } \mathrm{tw}(\Delta)} \text{holim } \Phi_\alpha & \xrightarrow{\text{proj.}} & \text{holim } \Phi_{\mathrm{id}_0} \end{array}$$

where id_0 is the identity morphism of $[0]$ in Δ. (The map from holim Φ to $\prod_\alpha$ holim Φ_α in this diagram has coordinates equal to the prolongation maps holim $\Phi \to$ holim Φ_α.) We are going to show that the two arrows in the lower row of diagram (1.2.3) are weak equivalences. For the one on the left it is a routine task. We can write

$$\begin{aligned} \operatorname*{holim}_{\alpha\colon[t]\to[u]} \text{holim } \Phi_\alpha &= \operatorname*{holim}_{\alpha\colon[t]\to[u]} \mathrm{Tot}\Big([r] \mapsto \prod_{\sigma_0 \ge \cdots \ge \sigma_r \ \in P_u} \lim(\Phi \circ \sigma_r\alpha_*)\Big) \\ &= \mathrm{Tot}\Big([r] \mapsto \operatorname*{holim}_{\alpha\colon[t]\to[u]} \prod_{\sigma_0 \ge \cdots \ge \sigma_r \ \in P_u} \lim(\Phi \circ \sigma_r\alpha_*)\Big) \end{aligned}$$

where

$$\operatorname*{holim}_{\alpha\colon[t]\to[u]} \prod_{\sigma_0\geq\cdots\geq\sigma_r\ \in P_u} \lim(\Phi\circ\sigma_r\alpha_*)$$

is the space of lifts as in the following commutative diagram:

(1.2.4)



(Note that N_rP_u is the set of singular u-simplices of the space $N_r\mathcal{P}(L)$, and so the bottom term in diagram (1.2.4) is the geometric realization of the singular simplicial set of the space $N_r\mathcal{P}(L)$.) But since the horizontal map in diagram (1.2.4) is a weak equivalence, the map from the section space $\Gamma_r(\Phi)$ of $E^!_r(\Phi)\to N_r\mathcal{P}(L)$ to

$$\operatorname*{holim}_{\alpha\colon[t]\to[u]} \prod_{\sigma_0\geq\cdots\geq\sigma_r\ \in P_u} \lim(\Phi\circ\sigma_r\alpha_*)$$

that it induces is a weak equivalence for every $r\geq 0$. It follows (by application of Tot) that the first arrow in the lower row of (1.2.3) is a weak equivalence.
For the other arrow in the lower row of diagram (1.2.3) it suffices to show that the functor

$$\big(\,\alpha\colon[t]\to[u]\,\big)\ \mapsto\ \text{holim}\ \Phi_\alpha$$

on $\mathrm{tw}(\Delta)$ takes all morphisms to weak equivalences. This reduces easily to the weaker statement where we only consider morphisms between objects of the form $\alpha\colon[0]\to[u]$ in $\mathrm{tw}(\Delta)$. For such an α it suffices to show that the map

$$\text{holim}\ \Phi_{\mathrm{id}_0}\to\text{holim}\ \Phi_\alpha$$

induced by the unique morphism from α to $\mathrm{id}_0\colon[0]\to[0]$ is a weak equivalence. Now [**5**, thm 6.12] can be applied. Then it only remains to show that the functor $\alpha^*\colon P_u\to P_0$ is homotopically terminal, i.e., that for every element z of P_0 the over category $P_{u,z}:=z/\alpha^*$ has a contractible classifying space. Note that $P_{u,z}$ is just the full sub-poset of P_u consisting of all σ in P_u such that $z\leq\sigma(w)$ in P_0, where w is the vertex $\alpha_*(0)$ of Δ^u.
For $z=(S,\rho)\in P_0$ and $\sigma\in P_{u,z}$ let $F(\sigma)$ be the space of *based* continuous maps $g\colon\Delta^u\to\mathrm{emb}(S,L)$ such that $g(x)$ takes S to $V_L(\sigma(x))$, for all $x\in\Delta^u$. (*Based* means that $g(w)$ is the inclusion $S\to L$.) Clearly $F(\sigma)$ is contractible and F is a covariant functor. So the projection

$$\text{hocolim}\ F\longrightarrow|N(P_{u,z})|$$

is a weak equivalence. But there is also a forgetful projection from hocolim F to the space Y of all based (continuous) maps $g\colon\Delta^u\to\mathrm{emb}(S,L)$. It is a fiber bundle

projection with contractible base space. Therefore it remains only to show that the fibers of this are contractible. Each fiber Y_g has the form $|N(P_{u,z,g})|$ where $P_{u,z,g}$ is the full sub-poset of $P_{u,z}$ consisting of the $\sigma \in P_{u,z}$ which in addition to the conditions for membership in $P_{u,z}$ satisfy $g(x)(S) \subset V_L(\sigma(x))$ for all $x \in \Delta^u$. It is easy to see that the poset $P_{u,z,g}$ satisfies a form of directedness and so $|N(P_{u,z,g})|$ is contractible. □

REMARK 1.2.2. For an open subset U of L, let $\mathcal{P}(L)|_U$ be the full topological sub-poset of $\mathcal{P}(L)$ consisting of all (S, ρ) such that $V_L(S, \rho)$ is contained in U. Let $\delta\mathcal{P}(L)|_U$ be the corresponding discrete poset. In the proof of theorem 1.1.1, we shall need a variant of lemma 1.2.1 which states that the comparison map

$$\operatorname*{holim}_{(S,\rho) \text{ in } \mathcal{P}(L)|_U} \Phi(S, \rho) \longrightarrow \operatorname*{holim}_{(S,\rho) \text{ in } \delta\mathcal{P}(L)|_U} \Phi(S, \rho)$$

is a weak equivalence. The proof is as for lemma 1.2.1.

1.3. Using manifold calculus. Let $\mathcal{O}(L)$ be the (discrete) poset of open subsets of L. In view of lemma 1.2.1, the following plan for a proof of theorem 1.1.1 looks promising. There is a contravariant functor

$$V \mapsto M \smallsetminus V \tag{1.3.1}$$

from $\mathcal{O}(L)$ to spaces. Manifold calculus as in [**11**] and [**8**] was created to help in understanding such functors. In particular, if a contravariant functor F from $\mathcal{O}(L)$ to spaces has some reasonable properties such as isotopy invariance and satisfies some approximate excision conditions, then manifold calculus has a formula

$$F(L) \xrightarrow{\simeq} \operatorname*{holim}_{U \in \bigcup_k \mathcal{O}k(L)} F(U) \ .$$

(Here $\mathcal{O}k(L) \subset \mathcal{O}(L)$ is the full sub-poset whose elements are the open subsets of L which are abstractly diffeomorphic to $\mathbb{R}^\ell \times S$ for some set S with $|S| \leq k$. Therefore

$$\bigcup_k \mathcal{O}k(L) \subset \mathcal{O}(L)$$

is the full sub-poset whose elements are the open subsets of L which are abstractly diffeomorphic to $\mathbb{R}^\ell \times S$ for some finite set S.) Using lemma 1.2.1, we should be able to work from there to arrive at theorem 1.1.1.

There is a small problem with this plan. The functor (1.3.1) does not have all the good properties required such as isotopy invariance. (Example: take L to be $S^2 = \mathbb{R}^2 \cup \infty$ and take M to be $S^m = \mathbb{R}^m \cup \infty$ for some $m \geq 2$. Let $V_1 \subset \mathbb{R}^2 \subset L$ be the union of the open rectangles $]2^{-i-1}, 2^{-i}[\times]0, 1[$ for $i = 0, 1, 2, \ldots$ and let $V_0 \subset \mathbb{R}^2 \subset L$ be the union of the open squares $]2^{-i-1}, 2^{-i}[\times]0, 2^{-i-1}[$ for $i = 0, 1, 2, \ldots$. The inclusion $V_0 \to V_1$ is an isotopy equivalence, but the induced homomorphism $\pi_{m-1}(M \smallsetminus V_1) \to \pi_{m-1}(M \smallsetminus V_0)$ is not surjective.) But that is easy to fix. We rectify (1.3.1) by setting

$$F(V) := \operatorname*{holim}_{C \subset V} M \smallsetminus C \tag{1.3.2}$$

for $V \in \mathcal{O}(L)$, where C runs through all compact subsets of V. Note that $F(L)$ has a forgetful projection map to $M \setminus L$ which is a weak equivalence, for if $V = L$ in (1.3.2), then there is a maximal choice for C which is $C = L$. Moreover, that map $F(L) \to M \setminus L$ has a preferred section which can be obtained by composing $M \setminus L \to \lim_C (M \setminus C) \to \operatorname{holim}_C (M \setminus C)$. The section is therefore also a weak equivalence. Now we need to show that F has reasonable properties such as isotopy invariance, and that it satisfies some approximate excision conditions.

LEMMA 1.3.1. *The functor F of* (1.3.2) *is* good. *That is to say:*

- *if $V_0 \subset V_1$ are open subsets of L such that the inclusion $V_0 \to V_1$ is abstractly isotopic to a diffeomorphism, then the map $F(V_1) \to F(V_0)$ induced by the inclusion is a weak homotopy equivalence;*
- *if $W \in \mathcal{O}(L)$ is a union of open subsets W_i where $i = 0, 1, 2, 3, \ldots$ and $W_i \subset W_{i+1}$, then the map from $F(W)$ to $\operatorname{holim}_i F(W_i)$ determined by the inclusions $W_i \to W$ is a weak equivalence.*

PROOF. The second of the two properties claimed is obvious from the definition of F. By contrast the first property is not easy to establish. Choose a sequence $(C_i)_{i \geq 0}$ of compact subsets of V_1 such that $C_i \subset C_{i+1}$ for all $i \geq 0$ and every compact subset of V_1 is contained in one of the C_i . Then the projection from $F(V_1)$ to the sequential homotopy limit

$$\operatorname{holim}_i M \setminus C_i$$

is a weak equivalence. Choose a smooth isotopy $(e_t \colon V_0 \to V_1)_{t \in [0,1]}$ such that $e_0 \colon V_0 \to V_1$ is the inclusion and $e_1 \colon V_0 \to V_1$ is a diffeomorphism. (See remark 1.3.4.) Let $C_{t,i} := e_t(e_1^{-1}(C_i))$ and note that C_i has just been renamed $C_{1,i}$. Now the projection from $F(V_0)$ to

$$\operatorname{holim}_i M \setminus C_{0,i}$$

is also a weak equivalence. Let Y_i be the space of continuous maps $w \colon [0,1] \to M$ such that $w(t) \notin C_{t,i}$ for all $t \in [0,1]$. By a straightforward application of Thom's isotopy extension theorem, the maps

$$M \setminus C_{0,i} \longleftarrow Y_i \longrightarrow M \setminus C_{1,i}$$

given by evaluation, $w \mapsto w(0)$ and $w \mapsto w(1)$, are homotopy equivalences. Therefore in the resulting diagram of sequential homotopy limits

$$\operatorname{holim}_i M \setminus C_{0,i} \longleftarrow \operatorname{holim}_i Y_i \longrightarrow \operatorname{holim}_i M \setminus C_{1,i}$$

the two arrows are also weak equivalences. Now choose a monotone injective function $\psi \colon \mathbb{N} \to \mathbb{N}$ such that $C_{1,\psi(i)}$ contains $C_{t,i}$ for all $t \in [0,1]$. This gives us a diagram

$$\begin{array}{ccccc}
F(V_1) & & \xrightarrow{\text{induced by incl.}} & & F(V_0) \\
\downarrow \simeq & & & & \downarrow \simeq \\
\operatorname{holim}_i M \setminus C_{1,i} & \xrightarrow[\simeq]{a} & \operatorname{holim}_i M \setminus C_{1,\psi(i)} & \xrightarrow{b} & \operatorname{holim}_i M \setminus C_{0,i} \\
& \nwarrow^{\simeq} & & \nearrow_{\simeq} & \\
& & \operatorname{holim}_i Y_i & &
\end{array}$$

where the vertical arrows and the arrow a are forgetful projections while the arrow b is induced by the inclusions $C_{0,i} \to C_{1,\psi(i)}$ for each $i \in \mathbb{N}$. The top square and the bottom triangle in the diagram are homotopy commutative; we leave the verification to the reader. Therefore b is a weak equivalence and then the top horizontal arrow is a weak equivalence. □

Proof of theorem 1.1.1. We start by showing that the functor F of (1.3.2) is analytic and by giving some excision estimates for it. Since we know already that F is good, it suffices to look into the following situation. Let P be a smooth compact codimension zero subobject (submanifold) of L and let $Q_0, \dots Q_k$ be pairwise disjoint compact codimension zero subobjects of $L \smallsetminus \mathrm{int}(P)$. See [**8**] for terminology. In more detail, this means that P is a compact smooth codimension zero submanifold with boundary of L but each Q_i is a compact smooth manifold with corners,

$$\partial Q_i = \partial_0 Q_i \cup \partial_1 Q_i$$

where $\partial_0 Q_i = Q_i \cap \partial P$ and $\partial_1 Q_i$ is the closure of $\partial Q_i \smallsetminus P$ in Q_i. There is the concept of handle index [**8**, §0] of Q_i. (*Important*: intuitively this is the handle index of Q_i relative to the subspace $\partial_0 Q_i$.) We assume that Q_i has handle index q_i . For $S \subset \{0, 1, \dots, k\}$ let $Q_S := P \cup \bigcup_{i \in S} Q_i$ and put

$$W_S := \mathrm{int}(Q_S)$$

(taking the interior in L). The commutative cube of spaces

$$S \mapsto F(W_S)$$

determines a map from $F(W_{\{0,1,\dots,k\}})$ to

$$\operatorname*{holim}_{S \subsetneq \{0,1,2,\dots,k\}} F(W_S)\,.$$

We need an estimate for the connectivity of that map, in terms of the dimensions m and ℓ, the number k and and the numbers q_i for $i = 0, 1, \dots, k$. Because of our special assumption we can work instead with the cube

$$S \;\mapsto\; G(W_S) := M \smallsetminus W_S$$

(contravariant in the variable S). That cube is a strongly cocartesian cube in Goodwillie's terminology [**7**]. The map from $G(W_{\{0,1,\dots,k\}})$ to $G(W_{\{0,1,\dots,k\} \smallsetminus \{i\}})$ is $(m - q_i - 1)$-connected. Therefore the excision estimates of [**6**] and [**7**, Thm.2.3] for such cubes apply here. Plugging these estimates into [**8**, Defn. 2.1] we deduce that F is $(m-2)$-analytic with excess 1 for manifold calculus purposes. More to the point, the comparison map

$$F(L) \longrightarrow \operatorname*{holim}_{U \in \bigcup_k \mathcal{O}k(L)} F(U)$$

is a weak equivalence and the comparison map

$$F(L) \longrightarrow \operatorname*{holim}_{U \in \mathcal{O}k(L)} F(U)$$

is $(1 + (k+1)(m - \ell - 2))$-connected by [**8**, Thm. 2.3]. —

Now we need to relate $\bigcup_k \mathcal{O}k(L)$ to $\delta\mathcal{P}(L)$. There is a full monomorphism of posets

$$\delta\mathcal{P}(L) \to \bigcup_k \mathcal{O}k(L)$$

which takes (S,ρ) to $V_L(S,\rho)$. This leads to a commutative diagram

$$\begin{array}{ccc} M \smallsetminus L & \longrightarrow & \underset{(S,\rho)\in\delta\mathcal{P}(L)}{\operatorname{holim}} M \smallsetminus V_L(S,\rho) \\ \downarrow \simeq & & \simeq \downarrow \\ F(L) & \longrightarrow & \underset{(S,\rho)\in\delta\mathcal{P}(L)}{\operatorname{holim}} F(V_L(S,\rho)) \\ \uparrow = & & \uparrow \\ F(L) & \xrightarrow{\simeq} & \underset{U\in \bigcup_k \mathcal{O}k(L)}{\operatorname{holim}} F(U) \end{array} \tag{1.3.3}$$

Since we want to know that the top horizontal arrow is a weak equivalence (keeping lemma 1.2.1 in mind), we ought to show that the lower right-hand vertical arrow, call it g, is a weak equivalence. By a standard argument from the theory of homotopy limits [**4**, thm.9.7] and with notation as in remark 1.2.2, the map g is a weak equivalence if for each $U \in \bigcup_k \mathcal{O}k(L)$ the map

$$g_U : F(U) \longrightarrow \underset{(S,\rho)\in\delta\mathcal{P}(L)|_U}{\operatorname{holim}} F(V_L(S,\rho))$$

induced by the inclusions $V_L(S,\rho) \to U$ is a weak equivalence. To show that this is the case, fix U and choose open $W \subset U$ such that $W = V_L(T,\sigma)$ for some (T,σ) in $\mathcal{P}(L)$ and the inclusion $W \to U$ induces a bijection on π_0. Then in the commutative diagram

$$\begin{array}{ccc} F(U) & \xrightarrow{g_U} & \underset{(S,\rho)\in\delta\mathcal{P}(L)|_U}{\operatorname{holim}} F(V_L(S,\rho)) \\ \downarrow r_1 & & \downarrow r_2 \\ F(W) & \xrightarrow{g_W} & \underset{(S,\rho)\in\delta\mathcal{P}(L)|_W}{\operatorname{holim}} F(V_L(S,\rho)) \end{array}$$

the map r_1 is clearly a weak equivalence. The map r_2 is a weak equivalence, too. Indeed by remark 1.2.2 it is allowed to replace $\delta\mathcal{P}(L)|_U$ by $\mathcal{P}(L)|_U$ and $\delta\mathcal{P}(L)|_W$ by $\mathcal{P}(L)|_W$, which means that in the right-hand column we can have homotopy limits in the style of definition 1.1.2. Then the verification amounts to seeing that a certain map between cosimplicial spaces is a degreewise equivalence. Finally the map g_W is a weak equivalence by a cofinality argument. (There is a maximal element in $\delta\mathcal{P}(L)|_W$.) Therefore g_U is a weak equivalence as claimed.

This takes care of the first part of theorem 1.1.1. The proof of the second part follows similar lines. □

Remark 1.3.2. The above proof of theorem 1.1.1 might suggest that the k-th Taylor approximation of the functor F of (1.3.2), in the sense of manifold calculus, can be obtained by post-composing F with the k-th Taylor approximation of the identity functor from spaces to spaces, in the sense of Goodwillie's homotopy functor calculus. Surprisingly, this is false. (It is also easy to see that it is false in the case $k = 1$.) A partial explanation is as follows. If G is a functor from spaces to spaces

which is polynomial of degree $\leq k$ in the sense of homotopy functor calculus, then GF is polynomial of degree $\leq k$ in the manifold calculus sense. This is due to the similarity of the definitions of *polynomial functor* in the two functor calculuses, and to a property of F which was emphasized in the proof above. But if G is a functor from spaces to spaces which is homogeneous of degree k in the sense of homotopy functor calculus, then GF need not be homogeneous of degree k in the sense of manifold calculus. This is due to obvious differences in the classification of homogeneous functors in the two functor calculuses.

REMARK 1.3.3. How useful, interesting or faithful is the map (1.1.2) when the codimension $m - \ell$ is less than 3 ? Here is a codimension 2 case which is not encouraging. Let $M = S^3$ and let L be a knot in S^3, your favorite knot, but not the unknot. Let $\mathbb{Z}_\infty$ be the Bousfield-Kan $\mathbb{Z}$-completion functor from spaces to spaces. It comes with a natural transformation $e\colon \mathrm{id} \to \mathbb{Z}_\infty$. For simply connected spaces X, the natural map $e\colon X \to \mathbb{Z}_\infty X$ is a weak homotopy equivalence; this is applicable when X is $\Phi(S,\rho)$ for some $(S,\rho) \in \mathcal{P}(L)$. But for $X = M \smallsetminus L$ the natural map $X \to \mathbb{Z}_\infty X$ is not a weak equivalence because $\mathbb{Z}_\infty(M \smallsetminus L) \simeq S^1$. (Instead it is a well-known map $M \smallsetminus L \to S^1$ which induces an isomorphism in ordinary integer homology.) We obtain a commutative diagram

$$\begin{array}{ccc} M \smallsetminus L & \xrightarrow{(1.1.2)} & \mathrm{holim}\ \Phi \\ \Big\downarrow{\scriptstyle e} & & \simeq \Big\downarrow{\scriptstyle e} \\ \mathbb{Z}_\infty(M \smallsetminus L) & \longrightarrow & \mathrm{holim}\ (\mathbb{Z}_\infty \circ \Phi) \end{array}$$

It follows that the map (1.1.2), top horizontal arrow in the diagram, factors up to homotopy through the notorious map $M \smallsetminus L \to S^1$. That seems to make (1.1.2) tragically un-faithful, in this codimension 2 example.

REMARK 1.3.4. In articles on manifold calculus, the meaning of *isotopy equivalence* is sometimes ambiguous. According to one definition, call it (a), a smooth codimension zero embedding $e\colon U \to V$ (of smooth manifolds with empty boundary) is an isotopy equivalence if and only if there exists an embedding $f\colon V \to U$ such that ef and fe are smoothly isotopic to the respective identity maps. According to another definition, call it (b), the embedding $e\colon U \to V$ is an isotopy equivalence if and only if it is isotopic (as a smooth embedding) to a diffeomorphism from U to V. We do not know whether definitions (a) and (b) are equivalent. Fortunately it is easy to see that, if a functor from $\mathcal{O}(L)$ to spaces takes isotopy equivalences as in definition (b) to weak equivalences, then it takes isotopy equivalences as in definition (a) to weak equivalences. (The converse is obvious.)

REMARK 1.3.5. Two slightly different views exist on what manifold calculus is about. In the older view laid out in [**11**] and [**8**], manifold calculus is about (some) contravariant functors from $\mathcal{O}(L)$ to spaces, where L is a fixed background manifold. In a more modern view, described for example in [**2**] though it was also heralded in [**1**], manifold calculus is about contravariant functors from a certain category Man_ℓ of *all* smooth ℓ-manifolds to spaces (for some ℓ). The morphisms in Man_ℓ are smooth embeddings between ℓ-manifolds. More precisely, the morphisms from L_0 to L_1 are organized into a space (or simplicial set), composition of morphisms is continuous (or is a simplicial map) etc., which means that Man_ℓ is *enriched* over the

category of spaces (or simplicial sets). Similarly the category of spaces is enriched over spaces (or simplicial sets), and the contravariant functors from Man_ℓ to spaces that we consider in manifold calculus should respect the enrichments.

A functor like $\mathrm{emb}(-,W)$ for a fixed smooth manifold W lives comfortably in both settings: the placeholder $-$ can be interpreted as an open subset of the fixed manifold L, or as an object of Man_ℓ. By contrast the functor F of (1.3.2) which we have used in proving theorem 1.1.1 seems to belong to the older setting; $F(V)$ makes sense only for open subsets V of L.

Does this mean that the modern reformulation of manifold calculus as in [**2**] has thrown out the baby with the bathwater? Nothing could be further from the truth. M.W. believes that most of the old manifold calculus can be subsumed in the new one as the branch concerned with contravariant functors G from Man_ℓ to spaces (preserving enrichment) which come equipped with a natural transformation γ to a representable functor

$$\mathrm{mor}_{\mathsf{Man}_\ell}(-,L) = \mathrm{emb}(-,L)$$

for fixed L in Man_ℓ. Such a pair (G,γ) gives rise to a contravariant functor G_γ from $\mathcal{O}(L)$ to spaces by

$$G_\gamma(V) := \mathrm{hofiber}[\gamma\colon G(V) \to \mathrm{emb}(V,L)]$$

for $V \in \mathcal{O}(L)$, where the homotopy fiber is taken over the base point of $\mathrm{emb}(V,L)$. The construction $(G,\gamma) \to G_\gamma$ should be seen as a transform, i.e., it is often reversible. In particular most of the contravariant functors from $\mathcal{O}(L)$ to spaces that we encounter in the old manifold calculus are weakly equivalent to G_γ for some G and $\gamma\colon G \to \mathrm{emb}(-,L)$. *Exercise*: confirm this for the functor F of (1.3.2). — In any case, the proposed subsuming of the old manifold calculus in the new one has not yet been carried out. That was one major reason for not using it here.

2. Occupants in the interior of a manifold

2.1. Formulation of the problem. Let M be a smooth compact manifold with boundary. We look for a homotopical description of ∂M in terms of the spaces $M \smallsetminus S$, where S runs through the finite subsets of $M \smallsetminus \partial M$. To make that more precise, choose a Riemannian metric on M. Then $M \smallsetminus \partial M$ also has a Riemannian metric and the topological poset $\mathcal{P}(M \smallsetminus \partial M)$ is defined as in section 1. Thus, elements of $\mathcal{P}(M \smallsetminus \partial M)$ are pairs (S,ρ) where S is a finite subset of $M \smallsetminus \partial M$ and ρ is a function from S to the positive reals such that, for each $s \in S$, the exponential map $\exp_s\colon T_s(M \smallsetminus \partial M) \to M \smallsetminus \partial M$ is defined and regular on the disk of radius $\rho(s)$ about the origin, and the images of these disks in $M \smallsetminus \partial M$ are pairwise disjoint. For $(S,\rho) \in \mathcal{P}(M \smallsetminus \partial M)$ let $V(S,\rho) \subset M \smallsetminus \partial M$ be the union of the open balls of radius $\rho(s)$ about points $s \in S$. Then $V(S,\rho)$ is homeomorphic to $\mathbb{R}^m \times S$. A contravariant functor Ψ from $\mathcal{P}(M \smallsetminus \partial M)$ to spaces is defined by

$$\Psi(S,\rho) = M \smallsetminus V(S,\rho). \tag{2.1.1}$$

There are maps

$$\partial M \longrightarrow \mathrm{holim}\ \Psi\ , \tag{2.1.2}$$

$$\partial M \longrightarrow \mathrm{holim}\ \Psi|_{\mathcal{P}j(M \smallsetminus \partial M)} \tag{2.1.3}$$

induced by the inclusions $\partial M \to M \smallsetminus V(S,\rho)$. The precise definition of the homotopy limits follows the pattern of definition 1.1.2.

THEOREM 2.1.1. *Suppose that M is the total space of a smooth disk bundle $p\colon M \to L$ of fiber dimension c on a smooth closed manifold L. If $c \geq 3$, then the map* (2.1.2) *is a weak equivalence and the map* (2.1.3) *is* $(1 + (j+1)(c-2))$*-connected.*

The expression *disk bundle* means just that: a smooth fiber bundle whose fibers are diffeomorphic to disks D^c of a fixed dimension c. It is not necessary to assume that p is the disk bundle associated with a smooth vector bundle on L.

2.2. The tube lemma. Theorem 2.1.1 will be proved by a reduction to theorem 1.1.1. The main idea for the reduction is in lemma 2.2.1 below. The lemma uses the notation of theorem 2.1.1, but we can allow a disk bundle $p\colon M \to L$ of any fiber dimension ≥ 0. We choose a Riemannian metric on L. Then in addition to the topological poset $\mathcal{P}(M \smallsetminus \partial M)$ there is the topological poset $\mathcal{P}(L)$, and there are some interactions between the two which will be explored. For $(S,\rho) \in \mathcal{P}(M \smallsetminus \partial M)$ the open set $V(S,\rho) \subset M \smallsetminus \partial M$ was defined just above. To be more consistent with section 1 we ought to write $V_{M\smallsetminus\partial M}(S,\rho)$, but that would be cumbersome. For $(S,\rho) \in \mathcal{P}(L)$ we still write $V_L(S,\rho) \subset L$ in the style of section 1.

LEMMA 2.2.1. *The map*

$$\operatorname*{hocolim}_{(S,\rho)\in\delta\mathcal{P}(L)} C_k(p^{-1}(V_L(S,\rho)) \smallsetminus \partial M) \longrightarrow C_k(M \smallsetminus \partial M) \tag{2.2.1}$$

determined by the inclusions $C_k(p^{-1}(V_L(S,\rho)) \smallsetminus \partial M) \to C_k(M \smallsetminus \partial M)$ is a weak equivalence.

PROOF. It suffices to show that (2.2.1) is a Serre microfibration with contractible fibers [**12**, Lemma 2.2]. Contractibility of the fibers is straightforward. The fiber over a configuration $T \in C_k(M \smallsetminus \partial M)$ is identified with the classifying space of the sub-poset $\mathcal{H}_T$ of $\delta\mathcal{P}(L)$ consisting of all (S,ρ) such that $p(T) \subset V_L(S,\rho)$. The poset $\mathcal{H}_T$ is the target of a homotopy initial functor from the poset of the negative integers.

It remains to show that (2.2.1) is a Serre microfibration. Here it is important to understand that although (2.2.1) and the projection to $|N\delta\mathcal{P}(L)|$ together determine an injective continuous map

$$\operatorname*{hocolim}_{(S,\rho)\in\delta\mathcal{P}(L)} C_k(p^{-1}(V_L(S,\rho)) \smallsetminus \partial M) \longrightarrow |N\delta\mathcal{P}(L)| \times C_k(M \smallsetminus \partial M), \tag{2.2.2}$$

that injective continuous map is not an embedding (homeomorphism onto the image). Here is a lengthy example to illustrate the phenomenon and the advantages that it has for us. Suppose for simplicity that $p\colon M \to L$ is an identity map, i.e., disk bundle with fiber dimension 0. Take two elements (S,ρ) and (T,σ) of $\delta\mathcal{P}(L)$ such that $(S,\rho) > (T,\sigma)$. The inequality $(S,\rho) > (T,\sigma)$ determines a nondegenerate 1-simplex in $N(\delta\mathcal{P}(L))$, and an injective map $\Delta^1 \to |N\delta\mathcal{P}(L)|$. View that as a path $w\colon [0,1] \to N(\delta\mathcal{P}(L))$, beginning at (T,σ) and ending at (S,ρ). Suppose that w has a lift $\bar{w}$ to a path in $\operatorname{hocolim}_{(S,\rho)} C_k(V_L(S,\rho))$. It is clear that the composition

$$[0,1] \xrightarrow{\bar{w}} \operatorname*{hocolim}_{(S,\rho)\in\delta\mathcal{P}(L)} C_k(V_L(S,\rho)) \longrightarrow C_k(L) \tag{2.2.3}$$

has the form $t \mapsto R_t$ where the configuration R_t is contained in $V(T,\sigma)$ if $0 \leq t < 1$ and in $V(S,\rho)$ when $t = 1$. But more careful reasoning shows that R_1 must be

contained in $V(T,\sigma)$, too. (If a hint is needed, see remark 2.2.2 below.) This is fortunate for us because it implies at once that a sufficiently small homotopy of the composition (2.2.3) can be lifted to a homotopy of $\bar{w}$ itself, as the Serre microfibration condition wants to have it. — Now we return to our business, which is to establish the Serre microfibration condition for the map (2.2.1). Let Z be a compact CW-space equipped with a map f to the source in (2.2.1). Using the injection (2.2.2), we may write $f = (f_1, f_2)$ where f_1 is a map with target $|N\delta\mathcal{P}(L)|$ and f_2 is a map with target $C_k(M \smallsetminus \partial M)$. Let

$$h\colon Z \times [0,1] \to C_k(M \smallsetminus \partial M)$$

be a homotopy such that $h_0 = f_2$. Note that $f_1(Z)$ is contained in a finite union of cells of $|N\delta\mathcal{P}(L)|$. For sufficiently small $\varepsilon > 0$, the map

$$H\colon Z \times [0,\varepsilon] \longrightarrow \operatorname*{hocolim}_{(S,\rho)} C_k(p^{-1}(V_L(S,\rho)) \smallsetminus \partial M)$$

defined by the formula $H = (f_1, h)$ using (2.2.2) is therefore continuous and well defined. (The digression just above was meant to prepare for that slightly counter-intuitive claim.) □

Remark 2.2.2. On the theme of counter-intuitive topological properties of homotopy colimits in the category of spaces, M.W. learned the following from Larry Taylor many years ago. Let U be a bounded open interval in $\mathbb{R}$. The mapping cylinder of the inclusion $U \to \mathbb{R}$ is not homeomorphic to a subspace of $\mathbb{R}^2$. It is not metrizable.

Corollary 2.2.3. *The map*

$$\operatorname*{hocolim}_{(S,\rho)\in\delta\mathcal{P}(L)} N_0\mathcal{P}(p^{-1}(V_L(S,\rho)) \smallsetminus \partial M) \longrightarrow N_0\mathcal{P}(M \smallsetminus \partial M)$$

determined by the inclusions $p^{-1}(V_L(S,\rho)) \smallsetminus \partial M \longrightarrow M \smallsetminus \partial M$ is a weak equivalence.

Proof. This is obtained from lemma 2.2.1 essentially by taking the disjoint union over all $k \geq 0$, noting that the hocolim respects disjoint unions. Perhaps it should be clarified that $\mathcal{P}(U)$, for an open subset U of $M \smallsetminus \partial M$, is defined or can be defined as the full topological sub-poset of $\mathcal{P}(M \smallsetminus \partial M)$ consisting of all elements (S,ρ) such that $V(S,\rho)$ is contained in U and has compact closure in U. □

Corollary 2.2.4. *For every $r \geq 0$ the map*

$$\operatorname*{hocolim}_{(S,\rho)\in\delta\mathcal{P}(L)} N_r\mathcal{P}(p^{-1}(V_L(S,\rho)) \smallsetminus \partial M) \longrightarrow N_r\mathcal{P}(M \smallsetminus \partial M)$$

determined by the inclusions $p^{-1}(V_L(S,\rho)) \smallsetminus \partial M \longrightarrow M \smallsetminus \partial M$ is a weak equivalence.

Proof. This is obtained from the previous corollary by noting that for open U in $M \smallsetminus \partial M$ there is a homotopy pullback square

$$\begin{array}{ccc} N_r\mathcal{P}(U) & \longrightarrow & N_r\mathcal{P}(M \smallsetminus \partial M) \\ \downarrow & & \downarrow \\ N_0\mathcal{P}(U) & \longrightarrow & N_0\mathcal{P}(M \smallsetminus \partial M) \end{array}$$

where the horizontal arrows are inclusions and the vertical arrows are given by the ultimate target operator, also known as 0-th vertex operator. (The square is also a strict pullback square, but this is less relevant for us.) □

PROOF OF THEOREM 2.1.1, FIRST PART. There is a commutative square

$$\begin{array}{ccc} \underset{(T,\sigma)\in\mathcal{P}(M\smallsetminus\partial M)}{\operatorname{holim}} \Psi(T,\sigma) & \longrightarrow & \underset{(S,\rho)\in\delta\mathcal{P}(L)}{\operatorname{holim}} \underset{\substack{(T,\sigma)\in\mathcal{P}(M\smallsetminus\partial M)\\ \text{cls. of } p\,(V(T,\sigma))\subset V_L(S,\rho)}}{\operatorname{holim}} \Psi(T,\sigma) \\ \uparrow & & \uparrow \\ \partial M & \longrightarrow & \underset{(S,\rho)\in\delta\mathcal{P}(L)}{\operatorname{holim}} \partial M \cup \big(M \smallsetminus p^{-1}(V_L(S,\rho))\big) \end{array}$$

By corollary 2.2.4 the top horizontal arrow, given by specialization, is a weak equivalence. By theorem 1.1.1 and lemma 1.2.1, the lower horizontal arrow is a weak equivalence. We want to know that the left-hand vertical arrow is a weak equivalence. So it suffices to show that the right-hand vertical arrow is a weak equivalence. For that it suffices to show that for fixed $(S,\rho) \in \delta\mathcal{P}(L)$ the map

$$\begin{array}{c} \underset{\substack{(T,\sigma)\in\mathcal{P}(M\smallsetminus\partial M)\\ \text{cls. of } p\,(V(T,\sigma))\subset V_L(S,\rho)}}{\operatorname{holim}} \Psi(T,\sigma) \\ \uparrow \\ \partial M \cup \big(M \smallsetminus p^{-1}(V_L(S,\rho))\big) \end{array}$$

induced by the inclusion of $\partial M \cup \big(M \smallsetminus p^{-1}(V_L(S,\rho))\big)$ in the various $\Psi(T,\sigma)$ is a weak equivalence. The target can also be written as

$$\underset{(T,\sigma)\in\mathcal{P}(U)}{\operatorname{holim}} \Psi(T,\sigma) \tag{2.2.4}$$

where U is the open subset $p^{-1}(V_L(S,\rho)) \smallsetminus \partial M$ of $M \smallsetminus \partial M$. We now make a few alterations to that expression, which turn out to be weak equivalences under $\partial M \cup \big(M \smallsetminus p^{-1}(V_L(S,\rho))\big)$.

(1) Replace $\mathcal{P}(U)$ by $\delta\mathcal{P}(U)$.
(2) Let F be the contravariant functor from $\mathcal{O}(U)$ to spaces taking $W \in \mathcal{O}(U)$ to $\operatorname{holim}_C(M \smallsetminus C)$, where C runs through the compact subsets of W. Replace $\Psi(T,\sigma)$ by $F(V(T,\sigma))$.
(3) After implementing (1) and (2), replace $\delta\mathcal{P}(U)$ by $\bigcup_k \mathcal{O}k(U)$ and replace $F(V(T,\sigma))$ for $(T,\sigma) \in \delta\mathcal{P}(U)$ by $F(W)$ for $W \in \bigcup_k \mathcal{O}k(U)$.

Alterations (1) and (3) can be justified by arguments which we have seen in section 1. Alteration (2) is justified because there is a comparison map from $\Psi(T,\sigma)$ to $F(V(T,\sigma))$ which is a weak equivalence. In this way, expression (2.2.4) turns into

$$\underset{W\in\bigcup_k \mathcal{O}k(U)}{\operatorname{holim}} F(W)\,. \tag{2.2.5}$$

But the poset $\bigcup_k \mathcal{O}k(U)$ has a maximal element, which is U itself. Therefore expression (2.2.5) can be replaced by $F(U)$. It is easy to see that the reference map from $\partial M \cup \big(M \smallsetminus p^{-1}(V_L(S,\rho))\big)$ to $F(U)$ is a weak equivalence. □

PROOF OF THEOREM 2.1.1, SECOND PART. Fix $j > 0$ as in (2.1.3). We need a modification of lemma 2.2.1. Let k be another integer such that $j \geq k \geq 0$. The modification states that the projection map

$$\operatorname*{hocolim}_{(S,\rho)\in\delta\mathcal{P}j(L)} C_k(p^{-1}(V_L(S,\rho)) \smallsetminus \partial M) \longrightarrow C_k(M \smallsetminus \partial M)$$

is a weak equivalence. The proof is exactly like the proof of lemma 2.2.1 itself: the map is again a Serre microfibration with contractible fibers. We need $j \geq k$ for the contractibility of the fibers. — There is a commutative square

$$\begin{array}{ccc}
\operatorname*{holim}_{(T,\sigma)\in\mathcal{P}j(M\smallsetminus\partial M)} \Psi(T,\sigma) & \longrightarrow & \operatorname*{holim}_{(S,\rho)\in\delta\mathcal{P}j(L)} \operatorname*{holim}_{\substack{(T,\sigma)\in\mathcal{P}j(M\smallsetminus\partial M)\\ \text{cls. of } p\,(V(T,\sigma))\subset V_L(S,\rho)}} \Psi(T,\sigma) \\
\uparrow & & \uparrow \\
\partial M & \longrightarrow & \operatorname*{holim}_{(S,\rho)\in\delta\mathcal{P}j(L)} \partial M \cup \big(M \smallsetminus p^{-1}(V_L(S,\rho))\big)
\end{array}$$

By a modification of corollary 2.2.4 which is a consequence of the modification of lemma 2.2.1 just formulated, the top horizontal arrow is a weak equivalence. By theorem 1.1.1 and lemma 1.2.1, the lower horizontal arrow is $(1 + (j + 1)(c - 2))$-connected. We want to know that the left-hand vertical arrow is $(1+(j+1)(c-2))$-connected. So it suffices to show that the right-hand vertical arrow is a weak equivalence. This can be verified as in the proof of the first half of theorem 2.1.1. □

3. Gates

This section generalizes the previous two. Consequently it has two slightly different themes.

3.1. Submanifold case. For the first theme, imagine a smooth manifold M with boundary and a neat smooth compact submanifold L, so that $\partial L \subset \partial M$. We look for a homotopical description of $M \smallsetminus L$ in terms of the spaces $M \smallsetminus S$, where S runs through the finite subsets of $L \smallsetminus \partial L$. In the case where ∂L and ∂M are empty, this is exactly the situation of section 1. Also, in the case where ∂L is empty but ∂M is nonempty, it is almost exactly the situation of section 1 because in such a case it makes no substantial difference if we delete ∂M from M.
For a more precise formulation we extend the definition of $\mathcal{P}(L)$ given in section 1 so that L is allowed to have a nonempty boundary. Choose a Riemannian metric on L. The elements of $\mathcal{P}(L)$ are going to be pairs (S, ρ) where S is a finite subset of $L \smallsetminus \partial L$ and ρ is a function from $S \sqcup \partial L$ to the positive reals, locally constant on ∂L and subject to a few more conditions.

- For each $s \in S$, the exponential map $\exp_s$ at s is defined and regular on the disk of radius $\rho(s)$ about the origin in T_sL.
- The (boundary-normal) exponential map is defined and regular on the set of all tangent vectors $v \in T_zL$ where $z \in \partial L$, where the vector v is inward perpendicular to $T_z\partial L$ and $|v| \leq \rho(z)$.
- The images in L of these disks and the image of this band under the exponential map(s) are pairwise disjoint.

For a pair (S,ρ) satisfying these conditions, let $V_L(S,\rho) \subset L$ be the union of the open balls of radius $\rho(s)$ about elements $s \in S$ and the open collar on ∂L determined the normal distance function $\rho_{|\partial L}$. Then $V_L(S,\rho)$ is diffeomorphic to $(\mathbb{R}^\ell \times S) \sqcup [0,1[\times \partial L$ and the inclusion of $M \smallsetminus V_L(S,\rho)$ in $M \smallsetminus S$ is a homotopy equivalence. The partial order on $\mathcal{P}(L)$ is defined so that $(S_0,\rho_0) \leq (S_1,\rho_1)$ if and only if $V_L(S_0,\rho_0) \subset V_L(S_1,\rho_1)$. In this partial order the boundary ∂L acts like a gate which allows occupants to leave.
The poset $\mathcal{P}(L)$ can also be viewed as a category. A contravariant functor Φ from $\mathcal{P}(L)$ to spaces is defined by

$$\Phi(S,\rho) = M \smallsetminus V_L(S,\rho). \tag{3.1.1}$$

There is a map

$$M \smallsetminus L \longrightarrow \operatorname{holim} \Phi\ , \tag{3.1.2}$$

determined by the inclusions $M \smallsetminus L \to \Phi(S,\rho)$ for $(S,\rho) \in \mathcal{P}(L)$. Also, let $\mathcal{P}j(L)$ be the subspace and full topological sub-poset of $\mathcal{P}(L)$ consisting of all (S,ρ) in $\mathcal{P}(L)$ that satisfy $|S| \leq j$. Then again there is a map

$$M \smallsetminus L \longrightarrow \operatorname{holim} \Phi|_{\mathcal{P}j(L)} \tag{3.1.3}$$

determined by the inclusions $M \smallsetminus L \to \Phi(S,\rho)$.

THEOREM 3.1.1. *If $m - \ell \geq 3$, then the map* (3.1.2) *is a weak homotopy equivalence and the map* (3.1.3) *is $(1 + (j+1)(m-\ell-2))$-connected.*

The proof of this is very similar to the proof of theorem 1.1.1 and the details are therefore omitted.

3.2. Absolute case. Our second topic is a generalization of theorem 2.1.1 to a situation with more complicated boundary conditions. Let M be a compact smooth manifold with boundary and corners in the boundary. In particular ∂M is the union of two codimension zero smooth submanifolds $\partial_0 M$ and $\partial_1 M$ that intersect in the corner set

$$\partial_0 M \cap \partial_1 M = \partial(\partial_0 M) = \partial(\partial_1 M).$$

We look for a homotopical description of $\partial_1 M$ in terms of the spaces $M \smallsetminus S$, where S runs through the finite subsets of $M \smallsetminus \partial M$.
Now $M \smallsetminus \partial_1 M$ is a smooth manifold with boundary $\partial_0 M \smallsetminus \partial_1 M$; both $M \smallsetminus \partial_1 M$ and its boundary can be noncompact. Again we choose a Riemannian metric on all of M. For simplicity we require it to be a product metric in a neighborhood of $\partial_1 M$, i.e., the product of a Riemannian metric on $\partial_1 M$ and a Riemannian metric on $[0,\varepsilon]$ for some $\epsilon > 0$. Then we can define a topological poset $\mathcal{P}(M \smallsetminus \partial_1 M)$ roughly as in section 2. The elements are pairs (S,ρ) where S is a finite subset of $M \smallsetminus \partial M = (M \smallsetminus \partial_1 M) \smallsetminus \partial(M \smallsetminus \partial_1 M)$ and ρ is a function from $S \sqcup (\partial_0 M \smallsetminus \partial_1 M)$ to the positive reals which is locally constant on $\partial_0 M \smallsetminus \partial_1 M$ and subject to the usual conditions. The usual conditions are, briefly stated: regularity of the exponential maps on the tangential disks and the tangential closed band defined by ρ, and pairwise disjointness of their images in $M \smallsetminus \partial_1 M$. The union of the open balls of radius $\rho(s)$ about elements $s \in S$ and of the (half-)open band determined by the normal distance function ρ on $\partial_0 M$ is denoted by $V(S,\rho)$. Therefore $V(S,\rho)$ is an open subset of $M \smallsetminus \partial_1 M$ diffeomorphic to $(\mathbb{R}^m \times S) \sqcup (\partial_0 M \smallsetminus \partial_1 M) \times [0,1[$.

A contravariant functor Ψ from $\mathcal{P}(M \smallsetminus \partial_1 M)$ to spaces is defined by

$$\Psi(S, \rho) = M \smallsetminus V(S, \rho). \tag{3.2.1}$$

There are maps

$$\partial_1 M \longrightarrow \text{holim } \Psi\ , \tag{3.2.2}$$

$$\partial_1 M \longrightarrow \text{holim } \Psi|_{\mathcal{P}j(M \smallsetminus \partial M)} \tag{3.2.3}$$

induced by the inclusions $\partial_1 M \to M \smallsetminus V(S, \rho)$.

THEOREM 3.2.1. *Suppose that $M \smallsetminus \partial_1 M$ has a neat compact smooth submanifold L of codimension $c \geq 3$ making M into a smooth thickening of $L \cup \partial_0 M$; see definition* 3.2.2 *below. Then the map* (3.2.2) *is a weak equivalence and the map* (3.2.3) *is $(1 + (j + 1)(c - 2))$-connected.*

DEFINITION 3.2.2. On the meaning of *smooth thickening* in theorem 3.2.1: the main points are that the inclusion of $\partial_0 M \cup L$ in M is a homotopy equivalence and that the inclusion of $\partial_1 M$ in $M \smallsetminus L$ is a homotopy equivalence.
In detail, we start with a smooth compact manifold L, another smooth compact manifold A (think $A = \partial_0 M$) and a smooth embedding $u\colon \partial L \to A$ which avoids the boundary of A. Then we can speak of $L \cup A$, the pushout of $L \leftarrow \partial L \to A$. Let $Q \subset L$ be a closed collar, so that $\partial L \subset Q$ and there is a diffeomorphism $Q \to \partial L \times [0, 1]$ extending the map $x \mapsto (x, 0)$ on ∂L. Let L_1 be the closure of $L \smallsetminus Q$ in L. To make a smooth thickening of $L \cup A$ we need in addition

- a smooth disk bundle $E \to L_1$ whose total space has dimension $\dim(A)+1$;
- an identification of $E|_{\partial L_1}$ with the normal disk bundle of the embedding $u\colon \partial L \to A$, over the evident diffeomorphism $\partial L_1 \cong \partial L$.

Then the pushout T of $A \times [0, 1] \leftarrow E|_{\partial L_1} \to E$ is defined, on the understanding that the left-hand arrow embeds $E|_{\partial L_1}$ into $A \times \{1\}$. Now T is a compact manifold with boundary, smooth with corners. The corner set has three disjoint parts: $\partial A \times \{0\}$, $\partial A \times \{1\}$ and $\partial(E|_{\partial L_1})$. Think of ∂T as the union of $\partial_0 T := A \times \{0\}$ and $\partial_1 T$, the closure of $\partial T \smallsetminus \partial_0 T$ in ∂T. The parts of the corner set not accounted for by $\partial_0 T \cap \partial_1 T$ should be subjected to smoothing. There is a copy of $L = Q \cup L_1$ contained in T. And of course there is also a copy of A contained in T, in the shape of $A \times \{0\}$. Any smooth manifold with corners which is diffeomorphic to such a T relative to $L \cup A = L \cup \partial_0 T$ can be called a smooth thickening of $L \cup A$ or of $L \cup \partial_0 T$.

EXAMPLE 3.2.3. In theorem 3.2.1, the manifold M can be the total space of a smooth disk bundle $M \to L$ of fiber dimension $c \geq 3$, where L is allowed to be compact smooth *with boundary*, and $M_0 := M|_{\partial L}$. The proof of theorem 3.2.1 in this case is a straightforward variation of the proof of theorem 2.1.1.

3.3. Something like engulfing. We turn to the proof of theorem 3.2.1. This is broken up into remarks, definitions, lemmas and even a corollary.

REMARK 3.3.1. The second part of theorem 3.2.1 (the high connectivity statement) implies the first part (the weak equivalence statement). This is easy to see if we use the definition of holim Ψ as a subspace of the space of maps from $|N\mathcal{P}(M \smallsetminus \partial_1 M)|$ to M. The inclusion of $|N\mathcal{P}j(M \smallsetminus \partial_1 M)|$ in $|N\mathcal{P}k(M \smallsetminus \partial_1 M)|$, for $k > j$, is a cofibration. It follows that the projection

$$\text{holim } \Psi|_{\mathcal{P}k(M \smallsetminus \partial_1 M)} \longrightarrow \text{holim } \Psi|_{\mathcal{P}j(M \smallsetminus \partial_1 M)}$$

is a fibration for $k > j$. Therefore the canonical inclusion

$$\text{holim } \Psi \;=\; \lim_j \text{ holim } \Psi|_{\mathcal{P}j(M\smallsetminus\partial_1 M)} \;\longrightarrow\; \text{holim}_j \text{ holim } \Psi|_{\mathcal{P}j(M\smallsetminus\partial_1 M)}$$

is a weak equivalence. The homotopy groups π_r of the right-hand side can be calculated as inverse limits

$$\lim_j \pi_r \left(\text{holim } \Psi|_{\mathcal{P}j(M\smallsetminus\partial_1 M)}\right).$$

The higher derived inverse limit $\lim^1$ does not contribute to this calculation because of the Mittag-Leffler criterion [**10**, ch.7,App.]. The criterion is applicable here because we are assuming the second part of theorem 3.2.1.

Remark 3.3.2. Since the validity of theorem 3.2.1 does not depend on the Riemannian metric which we select for M, we can choose a Riemannian metric with very convenient properties. We shall assume that it is a product metric near $\partial_0 M$ as well, i.e., a neighborhood of $\partial_0 M$ is isomorphic as a Riemannian manifold to a product $\partial_0 M \times [0,\varepsilon]$ where the interval $[0,\varepsilon]$ has the standard Riemannian metric, and the isomorphism takes $\partial_0 M \subset M$ to $\partial_0 M \times \{0\} \subset \partial_0 M \times [0,\varepsilon]$. (The product structure near $\partial_0 M$ is automatically compatible with the product structure near $\partial_1 M$ which we assumed earlier, so that $\partial\partial_0 M = \partial_0 M \cap \partial_1 M$ has a neighborhood in M which is isomorphic to $\partial\partial_0 M \times [0,\varepsilon] \times [0,\varepsilon]$ as a Riemannian manifold.) As a result we have a standard compact collar for $\partial_0 M$ (of width ε). We shall also assume that the intersection of L with that collar has the form $\partial L \times [0,\varepsilon]$ in the collar coordinates.
In addition we choose an open tubular neighborhood U of L such that the closure $\bar{U}$ of U in M is a smooth disk bundle over L. Also, the intersection of U with the standard collar on $\partial_0 M$ (of width ε) is required to have the form $\partial U \times [0,\varepsilon]$ in the collar coordinates. Here $\partial U \subset \partial_0 M$ is a tubular neighborhood of ∂L whose closure in $\partial_0 M$ is a smooth disk bundle over ∂L.

Definition 3.3.3. Let $\mathcal{C}_1$ be the full topological sub-poset of $\mathcal{P}j(M \smallsetminus \partial_1 M)$ consisting of the objects (S,ρ) such that the locally constant function $\rho|_{\partial_0 M}$ is $\leq \varepsilon$ everywhere and the disks of radius $\rho(s)$ about elements $s \in S$ are all contained in U, the specified tubular neighborhood of L. Write $\delta\mathcal{C}_1$ for the discrete variant.

Definition 3.3.4. Let $\mathcal{C}_0$ be the full topological sub-poset of $\mathcal{P}j(M \smallsetminus \partial_1 M)$ consisting of the objects (S,ρ) such that $\rho \leq \varepsilon/3j$, and the closure of $V(S,\rho)$ is contained in the union of U and the standard collar on $\partial_0 M$ of width $\varepsilon/3j$. Nota bene: for an object (S,ρ) of $\mathcal{C}_0$ it can happen that S is not contained in U. Write $\delta\mathcal{C}_0$ for the discrete variant.

Lemma 3.3.5. *For every element (S,ρ) of $\delta\mathcal{C}_0$ there is some element (T,σ) of $\delta\mathcal{C}_1$ such that $(T,\sigma) \geq (S,\rho)$ in $\mathcal{P}j(M \smallsetminus \partial_1 M)$. Indeed the sub-poset of $\delta\mathcal{C}_1$ consisting of the elements (T,σ) which satisfy $(T,\sigma) \geq (S,\rho)$ has a contractible classifying space.*

Proof. It is enough to note that there is a real number τ, strictly between $\varepsilon/3j$ and ε, such that the parallel hypersurface to $\partial_0 M$ in M at distance τ from $\partial_0 M$ has empty intersection with the closure of $V(S,\rho)$. This is due to our assumption $|S| \leq j$ and the smallness of the radii in the metric balls which are part of $V(S,\rho)$. □

COROLLARY 3.3.6. *There is a homotopy commutative diagram of the following shape,*

$$\begin{array}{ccc} \operatorname{holim}\, \Psi|_{\mathcal{P}j(M\smallsetminus\partial_1 M)} & \longrightarrow & \operatorname{holim}\, \Psi|_{\delta\mathcal{C}_1} \\ \downarrow & & \vdots \\ \operatorname{holim}\, \Psi|_{\delta\mathcal{C}_0} & \overset{\simeq}{\cdots\cdots\cdots\cdots\cdots} & Y \end{array}$$

where the solid arrows are given by restriction.

PROOF. Define Y to be

$$\operatorname*{holim}_{(S,\rho)\in\delta\mathcal{C}_0} \quad \operatorname*{holim}_{\substack{(S,\rho)\geq(S',\,\rho')\leq(T,\sigma)\\ (T,\sigma)\in\delta\mathcal{C}_1\\ (S',\,\rho')\in\delta\mathcal{C}_0}} \quad \Psi(S',\rho')\,.$$

The inner homotopy limit is taken over the subset $\mathcal{M}(S,\rho)$ of the product poset $\delta\mathcal{C}_0 \times \delta\mathcal{C}_1$ consisting of pairs $((S',\,\rho'),(T,\sigma))$ which satisfy $(S,\rho) \geq (S',\,\rho')$ and $(S',\,\rho') \leq (T,\sigma)$ in $\mathcal{P}j(M\smallsetminus\partial_1 M)$. We note that $\mathcal{M}(S,\rho)$ is covariantly functorial in $(S,\rho) \in \delta\mathcal{C}_0$, so that $(S,\rho) \leq (\bar{S},\bar{\rho})$ in $\delta\mathcal{C}_0$ implies $\mathcal{M}(S,\rho) \subset \mathcal{M}(\bar{S},\bar{\rho})$. This implies that the inner homotopy limit is a contravariant functor of the variable (S,ρ). — The two broken arrows are then fairly obvious and the homotopy commutativity of the resulting diagram is also clear. The horizontal broken arrow is a weak equivalence by lemma 3.3.5. More precisely, there are comparison maps

$$\Psi(S,\rho) \quad\longrightarrow\quad \operatorname*{holim}_{\substack{(S,\rho)\geq(S',\,\rho')\\ (S',\,\rho')\in\delta\mathcal{C}_0}} \quad \Psi(S',\rho') \quad\longrightarrow\quad \operatorname*{holim}_{\substack{(S,\rho)\geq(S',\,\rho')\leq(T,\sigma)\\ (T,\sigma)\in\delta\mathcal{C}_1\\ (S',\,\rho')\in\delta\mathcal{C}_0}} \quad \Psi(S',\rho').$$

the second of which is prolongation along a forgetful functor. The first of these is a weak equivalence because the indexing poset in the target expression has a terminal (maximal) element. The second one is a weak equivalence because the forgetful functor from $\mathcal{M}(S,\rho)$ to the poset $\{\,(S',\rho') \in \delta\mathcal{C}_0 \mid (S,\rho) \geq (S',\rho')\}$ is homotopy terminal [**5**] by lemma 3.3.5. Therefore the composition is a weak equivalence, and this is what we need to conclude that the lower horizontal arrow in the diagram is a weak equivalence. □

LEMMA 3.3.7. *The map* $\operatorname{holim}\, \Psi|_{\mathcal{P}j(M\smallsetminus\partial_1 M)} \longrightarrow \operatorname{holim}\, \Psi|_{\mathcal{C}_0}$ *given by restriction is a weak equivalence. The map* $\operatorname{holim}\, \Psi|_{\mathcal{C}_0} \longrightarrow \operatorname{holim}\, \Psi|_{\delta\mathcal{C}_0}$ *given by restriction is also a weak equivalence*

PROOF. For the first statement, observe that the inclusion map from $N_r\mathcal{C}_0$ to $N_r\mathcal{P}j(M \smallsetminus \partial_1 M)$ is a homotopy equivalence for every $r \geq 0$. The second statement can be proved like lemma 1.2.1. □

LEMMA 3.3.8. *The composition*

$$\partial_1 M \xrightarrow{(3.2.3)} \operatorname{holim}\, \Psi|_{\mathcal{P}j(M\smallsetminus\partial_1 M)} \longrightarrow \operatorname{holim}\, \Psi|_{\mathcal{C}_1}$$

is $(1 + (j+1)(c-2))$*-connected.*

PROOF. Following the lines of section 2, reduce to the claim that the map

$$\partial_1 M \longrightarrow \operatorname*{holim}_{(S,\rho)\in\mathcal{P}j(L)} M \smallsetminus V_L(S,\rho) \tag{3.3.1}$$

induced by the inclusions $\partial_1 M \to M \smallsetminus V_L(S,\rho)$ is so-and-so highly connected. (Following the lines of section 2 gives a zigzag of weak equivalences *under* $\partial_1 M$ relating holim $\Psi|_{\mathcal{C}_1}$ to the target in (3.3.1).) The map (3.3.1) can be written as a composition

$$\partial_1 M \hookrightarrow M \smallsetminus L \longrightarrow \operatorname*{holim}_{(S,\rho)\in\mathcal{P}j(L)} M \smallsetminus V_L(S,\rho).$$

The first arrow is a homotopy equivalence by our definition of *thickening* and the second arrow is so-and-so-highly connected by theorem 3.1.1. □

Conclusion of proof of theorem 3.2.1. By remark 3.3.1 we can concentrate on the high connectivity statement, for a fixed $j \geq 0$. The homotopy commutative square of corollary 3.3.6 can be enlarged to a diagram

$$\begin{array}{ccccc}
\partial_1 M & \xrightarrow{(3.2.3)} & \text{holim } \Psi|_{\mathcal{P}j(M\smallsetminus\partial_1 M)} & \longrightarrow & \text{holim } \Psi|_{\mathcal{C}_1} \\
 & & \downarrow\simeq & & \downarrow \\
 & & & & \text{holim } \Psi|_{\delta\mathcal{C}_1} \\
 & & \downarrow & & \downarrow \\
 & & \text{holim } \Psi|_{\delta\mathcal{C}_0} & \xrightarrow{\simeq} & Y
\end{array}$$

The weak equivalence labels in the diagram are justified by lemma 3.3.7 and corollary 3.3.6. By lemma 3.3.8, the composition of the two arrows in the top row is highly connected. It follows that the map (3.2.3) is split injective on homotopy groups or homotopy sets in a certain range. But the second arrow in the top row induces a split injection on all homotopy groups or homotopy sets, too, as the right-hand part of the diagram shows. □

4. Occupants near the gate

Imagine a smooth compact manifold M with corners in the boundary, so that $\partial M = \partial_0 M \cup \partial_1 M$ as in theorem 3.2.1. (There is no need to assume here that M is a thickening as in definition 3.2.2.) One might ask for a homotopical description of the corner set $\partial\partial_1 M = \partial\partial_0 M = \partial_0 M \cap \partial_1 M$ in terms of the spaces $M \smallsetminus S$, where S runs through the finite subsets of $M \smallsetminus \partial M$ which are close to $\partial_0 M$. We only make a few steps towards such a description. The associated definitions turn out to be useful elsewhere [**13**].

4.1. The situation near the gate. Let $\mathcal{P}(M \smallsetminus \partial_1 M)$ be defined as in theorem 3.2.1. Form the twisted arrow poset $\mathrm{tw}(\mathcal{P}(M \smallsetminus \partial_1 M))$. (This is a special case of the twisted arrow category construction which was mentioned in the proof of lemma 1.2.1.) There is a contravariant functor Θ from $\mathrm{tw}(\mathcal{P}(M \smallsetminus \partial_1 M)$ to spaces given by

$$\Theta((S,\rho) \leq (T,\sigma)) := \big(\text{closure in } M \text{ of the collar part of } V(T,\sigma)\big) \smallsetminus V(S,\rho).$$

This comes with a natural inclusion map $\partial\partial_1 M \to \Theta\big((S,\rho) \leq (T,\sigma)\big)$ which in turn induces a map

$$\partial\partial_1 M \longrightarrow \operatorname{holim} \Theta\,. \tag{4.1.1}$$

There is a natural transformation $\Theta \to \Psi \circ F_s$ given by inclusion, where

$$F_s : \mathrm{tw}(\mathcal{P}(M \smallsetminus \partial_1 M)) \longrightarrow \mathcal{P}(M \smallsetminus \partial_1 M)$$

is the functor *source*. In more detail, for an object $(S,\rho) \leq (T,\sigma)$ of the poset $\mathrm{tw}(\mathcal{P}(M \smallsetminus \partial_1 M))$ the space $\Theta((S,\rho) \leq (T,\sigma))$ is obviously contained in $\Psi(S,\rho) = M \smallsetminus V(S,\rho)$. Using this we get a commutative diagram

$$\begin{array}{ccccc} \partial\partial_1 M & & \xrightarrow{\text{inclusion}} & & \partial_1 M \\ \downarrow{\scriptstyle (4.1.1)} & & & & \downarrow{\scriptstyle (3.2.2)} \\ \mathrm{holim}\ \Theta & \longrightarrow & \mathrm{holim}\ \Psi \circ F_s & \longleftarrow & \mathrm{holim}\ \Psi \end{array} \tag{4.1.2}$$

LEMMA 4.1.1. *The map from* holim Ψ *to* holim $\Psi \circ F_s$ *in* (4.1.2) *is a weak equivalence.*

PROOF. Recall that holim Ψ was defined as $\mathrm{Tot}(X)$ for a certain cosimplicial space X. Namely, X_r is the space of sections of the fiber bundle on $N_r\mathcal{P}(M \smallsetminus \partial M)$ whose fiber over $((S_0,\rho_0) \geq \cdots \geq (S_r,\rho_r))$ is $M \smallsetminus V(S_r,\rho_r)$. Let $\beta\colon \Delta \to \Delta$ be the functor $[n] \mapsto [2n+1]$. More precisely, Δ is the category of totally ordered nonempty finite sets and order-preserving maps, or the equivalent full subcategory with objects $[n]$ for $n \geq 0$, and β is the functor which takes a totally ordered nonempty finite set S to $S \sqcup S^{\mathrm{op}}$ (with the total ordering where $a < b$ whenever $a \in S \subset S \sqcup S^{\mathrm{op}}$ and $b \in S^{\mathrm{op}} \subset S \sqcup S^{\mathrm{op}}$). The inclusions $S \to S \sqcup S^{\mathrm{op}}$ define a natural transformation $e\colon \mathrm{id} \to \beta$. The cosimplicial space $X \circ \beta$ is Reedy fibrant, by the same argument which we used to show that X is Reedy fibrant. We need to show that the map $e_*\colon \mathrm{Tot}(X) \to \mathrm{Tot}(X \circ \beta)$ is a weak equivalence. Since both X and $X \circ \beta$ are Reedy fibrant, we can use the easier variant Tots of Tot where only the (co)face operators are used; but we continue to view X and $X \circ \beta$ as cosimplicial spaces. In this setting a more general statement can be made: *if Y is any cosimplicial space, then $e_*\colon \mathrm{Tots}(Y) \to \mathrm{Tots}(Y \circ \beta)$ is a weak equivalence.* To show this, we use standard resolution procedures and note that Tots preserves degreewise weak equivalences. Therefore we may assume that Y is a homotopy inverse limit of cosimplicial spaces having the form

$$[r] \mapsto \mathrm{map}(\hom_\Delta([r],[t]), Z) \tag{4.1.3}$$

for some $[t]$ in Δ and a space Z. Since Tots commutes with such homotopy inverse limits, it suffices to show that $e_*\colon \mathrm{Tots}(Y) \to \mathrm{Tots}(Y \circ \beta)$ is a weak equivalence when Y has the form (4.1.3). In that case $\mathrm{Tots}(Y)$ is just the space of maps from the geometric realization of the semi-simplicial set

$$[r] \mapsto \hom_\Delta([r],[t]) \tag{4.1.4}$$

to Z. Similarly $\mathrm{Tots}(Y \circ \beta)$ is the space of maps from the geometric realization of the semi-simplicial set

$$[r] \mapsto \hom_\Delta([2r+1],[t]) \tag{4.1.5}$$

to Z. Now it is enough to show that the geometric realizations of (4.1.4) and (4.1.5) are both contractible. For that we may pretend or observe that both (4.1.4) and (4.1.5) are actually simplicial sets and realize them as such. The result is in one case a standard geometric t-simplex. In the other case it is an edgewise subdivided t-simplex. □

4.2. A locality statement. We introduce another topological poset $\mathcal{Q} = \mathcal{Q}(M \smallsetminus \partial_1 M)$ which is closely related to $\mathrm{tw}(\mathcal{P}) = \mathrm{tw}(\mathcal{P}(M \smallsetminus \partial_1 M))$. Intuitively, $\mathcal{Q}$ is the quotient poset obtained from $\mathrm{tw}(\mathcal{P})$ by forcing a morphism in $\mathrm{tw}(\mathcal{P})$ to be an equality (in $\mathcal{Q}$) if the functor Θ takes it to an identity map of spaces. More formally, an element of $\mathcal{Q}$ is an element

$$\big((S,\rho) \leq (T,\sigma)\big)$$

of $\mathrm{tw}(\mathcal{P})$ where $T = \emptyset$, so that $V(T,\sigma)$ is nothing but a collar. For two such pairs

$$x = \big((S,\rho) \leq (T,\sigma)\big), \qquad x' = \big((S',\rho') \leq (T',\sigma')\big)$$

where $T = T' = \emptyset$, we say that $x \leq x'$ if $\Theta(x') \subset \Theta(x)$. As a result, $\mathcal{Q}$ is not a sub-poset of $\mathrm{tw}(\mathcal{P})$. Instead there is a functor (continuous map of posets) K from $\mathrm{tw}(\mathcal{P})$ to $\mathcal{Q}$ which takes

$$\big((S,\rho) \leq (T,\sigma)\big)$$

in $\mathrm{tw}(\mathcal{P})$ to $\big((S_0,\rho_0) \leq (T_0,\sigma_0)\big)$ where $T_0 = \emptyset$ and σ_0 is the appropriate restriction of σ, while S_0 is the part of S which is contained in the collar part of $V(T,\sigma)$ and ρ_0 is the appropriate restriction of ρ. By construction we have

$$\Theta = \Theta_1 \circ K$$

for a functor Θ_1 from $\mathcal{Q}$ to spaces. This gives us a prolongation map

$$\mathrm{holim}\ \Theta_1 \longrightarrow \mathrm{holim}\ \Theta_1 \circ K = \mathrm{holim}\ \Theta$$

and leads to a commutative diagram

$$\begin{array}{ccccccc} \partial\partial_1 M & & \xrightarrow{\qquad\qquad\text{inclusion}\qquad\qquad} & & \partial_1 M & & \\ \downarrow & & & & \downarrow & & (3.2.2) \\ \mathrm{holim}\ \Theta_1 & \longrightarrow & \mathrm{holim}\ \Theta & \longrightarrow & \mathrm{holim}\ \Psi \circ F_s & \xleftarrow{\simeq} & \mathrm{holim}\ \Psi \end{array} \tag{4.2.1}$$

which refines diagram (4.1.2). Namely, deletion of the vertex holim Θ_1 and composition of the two incident arrows recovers diagram (4.1.2).

In an older version of this article, it was claimed that the map from holim Θ_1 to holim Θ in diagram (4.2.1) is always a weak equivalence. The proof had many flaws. (If a more realistic proof of this comes to light, it might go into the next edition of [**13**].) There was also a conjecture which is best restated as saying that the map from $\partial\partial_1 M$ to holim Θ_1 in diagram (4.2.1) is a weak equivalence under some fairly restrictive conditions on $\partial_0 M$.

In an effort to practice restraint we finish with the following easy but important observation: the left-hand column in diagram (4.2.1) depends only on an arbitrarily small open neighborhood U of $\partial_0 M$ in M. To make a precise statement, let $\partial_1 U = U \cap \partial_1 M$ and $\partial_0 U = \partial_0 M$ and define $\mathcal{P}(U \smallsetminus \partial_1 U)$ by analogy with $\mathcal{P}(M \smallsetminus \partial_1 M)$. Alternatively define it as a full topological sub-poset of $\mathcal{P}(M \smallsetminus \partial_1 M)$, consisting of the objects (S,ρ) for which the closure of $V(S,\rho)$ in M is contained in U. Let $\Theta_{1,U}$ be the restriction of Θ_1 to $\mathrm{tw}(\mathcal{P}(U \smallsetminus \partial_1 U))$.

Proposition 4.2.1. *The forgetful projection* holim $\Theta_1 \longrightarrow$ holim $\Theta_{1,U}$ *is a weak equivalence.*

PROOF. The inclusion of the nerve of $\mathcal{Q}_U = \mathcal{Q}(U \smallsetminus \partial_1 U)$ in the nerve of $\mathcal{Q} = \mathcal{Q}(M \smallsetminus \partial_1 M)$ is a degreewise weak equivalence of simplicial spaces. □

* * *

Remark. An earlier version of this paper was circulated with M.W. as the sole author. But S.T. found so many inaccuracies in it that we decided to call the revised version a joint paper by S.T. and M.W.

References

1. G. Arone and V. Turchin, *On the rational homology of high-dimensional analogues of spaces of long knots*, Geom. Topol. 18 (2014), 1261-1322.
2. P. Boavida de Brito and M.S. Weiss, *Manifold calculus and homotopy sheaves*, Homology Homotopy Appl. 15 (2013), 361–383.
3. A Bousfield and D Kan, *Homotopy limits, completions and localizations*, Lecture Notes in Math. vol. 304, Springer-Verlag, Berlin-NewYork, 1972.
4. W. Dwyer and D. M. Kan, *A classification theorem for diagrams of simplicial sets*, Topology 23 (1984), 139–155.
5. D. Dugger, *A primer on homotopy colimits*, http://pages.uoregon.edu/ddugger/
6. G. Ellis and R. Steiner, *Higher dimensional crossed modules and the homotopy groups of* $(n+1)$*-ads*, J.Pure Appl. Algebra 46 (1987), 117-136.
7. T. G. Goodwillie, *Calculus II: Analytic Functors*, K-Theory 5 (1992), 295–332.
8. T. G. Goodwillie and M. S. Weiss, *Embeddings from the point of view of immersion theory, Part II*, Geometry and Topology 3 (1999), 103–118.
9. S.-T. Hu, *Theory of Retracts*, Wayne State University Press, 1965.
10. R. Switzer, *Algebraic Topology – Homotopy and Homology*, Grundlehren series 212, Springer-Verlag, 1975.
11. M. S. Weiss, *Embeddings from the point of view of immersion theory, Part I*, Geometry and Topology 3 (1999), 67–101.
12. M. S. Weiss, *What does the classifying space of a category classify ?*, Homology Homotopy Appl. 7 (2005), 185–195.
13. M. S. Weiss, *Configuration categories and homotopy automorphisms*, arXiv:1503.00501.

MATH. INSTITUT, UNIVERSITÄT MÜNSTER, 48149 MÜNSTER, EINSTEINSTRASSE 62, GERMANY
E-mail address: s_till05@uni-muenster.de
E-mail address: m.weiss@uni-muenster.de